

		Contraceptive prevalence rate	Skilled attendant at delivery	C-section rate	Antenatal care (4+ visits)	Early initiation of breastfeeding	Postnatal care for babies born at home	Postnatal care for all babies*	Family planning needs satisfied	Improved water sources	DPT3 vaccine	Measles vaccine	BCG vaccine	Vitamin A administration	ORT and continued feeding	Care seeking for pneumonia	Insecticide-treated net use by children	Co-coverage (6 or more)	Coverage gap	Mean coverage
Child's sex	Male	NA	76,0%	4,3%	46,6%	50,4%	NO	NO	NA	NA	46,1%	62,7%	74,8%	90,3%	45,0%	44,6%	NO	7,1%	NA	NA
	Female	NA	73,9%	3,9%	49,7%	50,4%	NO	NO	NA	NA	46,6%	65,0%	70,2%	89,3%	39,8%	39,1%	NO	6,4%	NA	NA
Region of residence	Urban	27,0%	91,8%	4,9%	54,2%	49,9%	NO	NO	52,0%	73,2%	60,8%	74,3%	85,5%	94,3%	40,6%	44,9%	NO	11,3%	NA	NA
	Rural	16,4%	64,1%	3,5%	44,0%	50,7%	NO	NO	40,6%	8,6%	36,9%	57,2%	64,2%	85,6%	43,4%	40,3%	NO	3,8%	NA	NA
Wealth quintiles	Q1	14,3%	60,2%	2,4%	42,9%	47,6%	NO	NO	36,7%	6,8%	28,8%	52,4%	62,6%	86,1%	38,7%	39,3%	NO	3,4%	57,4%	42,6%
	Q2	15,1%	63,1%	3,3%	45,1%	49,4%	NO	NO	39,1%	7,7%	37,1%	56,6%	63,5%	84,0%	41,0%	43,0%	NO	3,4%	54,0%	46,0%
	Q3	16,6%	73,9%	3,4%	46,3%	52,9%	NO	NO	40,0%	16,5%	45,4%	62,3%	63,8%	90,7%	47,4%	36,4%	NO	4,2%	50,9%	49,1%
	Q4	22,2%	85,9%	6,0%	45,2%	53,6%	NO	NO	46,8%	53,6%	53,4%	67,0%	83,2%	88,7%	43,9%	42,6%	NO	7,8%	45,0%	55,0%
	Q5	38,6%	98,3%	5,8%	64,1%	48,0%	NO	NO	62,0%	96,7%	73,9%	86,3%	95,5%	97,2%	39,8%	53,9%	NO	17,3%	31,9%	68,1%
Maternal education	A	11,8%	60,6%	4,8%	37,3%	50,0%	NO	NO	35,7%	NO	34,9%	49,9%	57,1%	89,7%	32,5%	36,1%	NO	2,7%	NA	NA
	B	17,4%	68,6%	3,3%	43,5%	52,0%	NO	NO	42,5%	NO	38,8%	59,1%	65,4%	86,8%	44,0%	40,2%	NO	4,5%	NA	NA
	C	26,0%	85,5%	4,0%	55,0%	50,0%	NO	NO	47,6%	NO	53,5%	71,3%	83,0%	90,6%	46,5%	42,9%	NO	9,3%	NA	NA
	D	38,6%	98,4%	5,8%	68,3%	45,9%	NO	NO	68,7%	NO	79,0%	90,7%	96,7%	96,9%	42,6%	72,5%	NO	16,6%	NA	NA
Country region	kinshasa	41,9%	97,5%	6,8%	69,0%	46,8%	NO	NO	64,0%	93,2%	82,6%	92,7%	94,8%	99,3%	47,8%	61,3%	NO	18,8%	NA	NA
	bas-congo	39,5%	93,1%	3,8%	50,4%	51,8%	NO	NO	63,5%	30,1%	83,5%	88,0%	92,1%	92,3%	46,8%	52,9%	NO	11,7%	NA	NA
	bandundu	26,2%	71,1%	2,5%	53,2%	48,5%	NO	NO	51,3%	4,8%	62,7%	79,0%	81,2%	93,7%	39,8%	42,4%	NO	6,5%	NA	NA
	equateur	16,0%	51,8%	1,2%	48,7%	51,6%	NO	NO	35,0%	9,5%	23,4%	42,4%	56,0%	87,0%	47,2%	34,3%	NO	0,9%	NA	NA
	orientale	11,8%	69,4%	2,7%	47,4%	43,5%	NO	NO	34,8%	18,6%	26,0%	50,4%	57,5%	81,8%	44,5%	47,0%	NO	5,0%	NA	NA
	nord-kivu	23,2%	88,9%	15,4%	50,3%	62,0%	NO	NO	41,6%	40,5%	84,8%	86,7%	94,5%	81,4%	50,4%	58,1%	NO	8,1%	NA	NA
	maniema	17,3%	70,5%	2,8%	44,4%	48,3%	NO	NO	45,1%	21,3%	18,7%	45,0%	55,7%	88,7%	46,4%	50,9%	NO	4,0%	NA	NA
	sud-kivu	13,8%	86,7%	10,8%	22,4%	45,4%	NO	NO	30,7%	44,4%	54,8%	85,0%	89,0%	97,8%	28,0%	42,3%	NO	6,7%	NA	NA
	katanga	19,9%	70,8%	1,8%	37,4%	32,8%	NO	NO	47,9%	60,5%	39,9%	52,6%	61,8%	84,2%	49,1%	43,9%	NO	6,0%	NA	NA
kasai orientale	11,3%	76,3%	3,5%	44,2%	57,7%	NO	NO	34,3%	57,3%	38,8%	59,6%	70,8%	86,2%	43,3%	30,3%	NO	6,9%	NA	NA	
kasai occidental	14,2%	79,7%	5,4%	43,7%	68,4%	NO	NO	38,7%	13,0%	27,8%	47,6%	68,4%	85,1%	32,0%	42,3%	NO	3,3%	NA	NA	
Overall		20,6%	74,9%	4,1%	48,2%	50,4%	NO	NO	45,9%	36,3%	46,3%	63,8%	72,5%	89,8%	42,3%	41,9%	NO	6,8%	48,3%	51,7%
Q5/Q1 ratio		2,70	1,63	2,38	1,49	1,01	NO	NO	1,69	14,28	2,57	1,65	1,53	1,13	1,03	NO	5,09	0,56	1,60	
Q5-Q1 difference		24,3	38,1	3,4	21,1	0,4	NO	NO	25,3	90,0	45,1	33,9	32,9	11,1	1,0	14,6	NO	13,9	-25,5	25,5
Slope index		27,8	47,0	4,6	19,1	3,3	NO	NO	30,8	107,9	52,1	37,5	41,0	13,1	3,2	12,3	NO	14,7	NA	NA
Concentration index		20,1	10,4	20,0	6,6	1,0	NO	NO	11,4	51,2	18,1	10,4	9,4	2,5	1,2	4,5	NO	37,0	NA	NA




Fig 1. Mean coverage of selected interventions and corresponding coverage gap (how much is needed to reach universal coverage), by wealth quintile.

	Antenatal care (4+ visits)	Skilled attendant at delivery	Care seeking for pneumonia	ORT and continued feeding	BCG vaccine	DPT3 vaccine	Measles vaccine	Family planning needs satisfied
Poorest quintile	42,9	60,2	39,3	38,7	62,6	28,8	52,4	36,7
Richest quintile	21,1	38,1	14,6	1,0	32,9	45,1	33,9	25,3
Richest quintile	64,1	98,3	53,9	39,8	95,5	73,9	86,3	62,0




Fig 2. Coverage levels in the poorest and richest quintiles for selected interventions along the continuum of care.

Co-coverage interval	Wealth quintile				
	Q1	Q2	Q3	Q4	Q5
0	0,199	0,209	0,100	0,047	0,000
1	0,275	0,247	0,231	0,136	0,011
2	0,225	0,287	0,335	0,283	0,198
3	0,156	0,123	0,157	0,254	0,323
4	0,076	0,065	0,092	0,140	0,222
5	0,035	0,035	0,041	0,062	0,073
6	0,030	0,025	0,033	0,058	0,091
7	0,004	0,008	0,006	0,017	0,067
8	0,000	0,000	0,003	0,003	0,014
9					
Total	1,000	1,000	1,000	1,000	1,000




Fig 3. Percentage of children aged 1-4 years according to the number of key child-survival interventions received, by wealth quintile.