

# Togo (MICS 2006)

Table - Coverage of Countdown indicators at national level and by key equity stratifiers.

	Family planning needs satisfied	Contraceptive prevalence rate	Antenatal care (1+ visits, skilled provider)	Antenatal care (4+ visits, any provider)	Skilled attendant at delivery	C-section rate	Early initiation of breastfeeding	Postnatal care for babies born at home	Postnatal care for all babies	BCG vaccine	DTP3 vaccine	Measles vaccine	Vitamin A in last 6 mos.	ORT and continued feeding	Care seeking for pneumonia	ITN use by children	Improved drinking water source	Co-coverage (6 or more interventions)	Composite coverage index	
<b>National</b>	29.3	16.8	82.4	NO	51.9	NO	35.8	NO	NO	88.0	63.9	63.1	39.3	37.5	22.7	38.5	60.6	16.0	49.1	
<b>Wealth quintiles</b>																				
Q1	21.7	12.1	67.5	NO	24.3	NO	28.8	NO	NO	84.1	58.0	57.2	35.4	33.6	18.4	40.5	31.1	9.7	39.5	
Q2	27.4	15.5	72.4	NO	22.9	NO	28.5	NO	NO	79.4	55.7	55.7	39.8	36.8	24.2	40.6	35.8	11.2	41.8	
Q3	29.6	17.0	82.9	NO	44.9	NO	42.6	NO	NO	87.4	60.8	60.7	39.0	34.7	6.1	41.3	54.1	13.0	45.4	
Q4	30.8	19.3	92.2	NO	79.1	NO	40.4	NO	NO	95.8	70.5	70.1	39.9	33.6	33.3	34.6	78.8	23.7	56.7	
Q5	35.8	19.4	97.7	NO	88.8	NO	40.0	NO	NO	93.9	74.8	72.1	42.6	56.4	27.7	35.4	94.4	23.2	62.5	
<b>Equity indicators</b>																				
Q5/Q1 ratio	1.7	1.6	1.4	NO	3.7	NO	1.4	NO	NO	1.1	1.3	1.3	1.2	1.7	1.5	0.9	3.0	2.4	1.6	
Q5-Q1 difference	14.1	7.3	30.1	NO	64.5	NO	11.2	NO	NO	9.7	16.8	14.9	7.2	22.8	9.3	-5.1	63.3	13.5	23.0	
Concentration index	8.6	8.9	8.2	NO	29.9	NO	8.2	NO	NO	3.7	6.6	6.1	3.3	5.7	9.4	-3.5	23.4	20.8	9.9	
SII	15.5	9.1	41.7	NO	81.5	NO	17.4	NO	NO	19.1	24.3	22.3	7.5	14.1	13.9	-8.1	76.9	20.1	30.4	
<b>Maternal education</b>																				
none	21.7	11.9	74.0	NO	36.4	NO	33.6	NO	NO	79.4	51.3	50.0	35.4	39.6	20.7	38.9	NA	10.2	41.3	
primary	32.3	18.8	87.8	NO	57.8	NO	36.9	NO	NO	93.2	70.7	68.7	42.8	30.4	23.3	37.1	NA	19.7	51.9	
secondary +	42.2	26.7	93.0	NO	78.8	NO	38.9	NO	NO	97.9	78.9	82.0	43.9	49.4	25.5	40.4	NA	25.5	62.5	
<b>Child's sex</b>																				
male	NA	NA	ND	NO	ND	NO	ND	NO	NO	89.0	65.3	60.9	38.0	37.0	22.7	40.3	NA	15.8	NA	
female	NA	NA	ND	NO	ND	NO	ND	NO	NO	87.1	62.6	65.2	40.6	38.0	22.8	36.6	NA	16.2	NA	
<b>Area of residence</b>																				
urban	30.5	17.5	94.0	NO	85.2	NO	39.5	NO	NO	93.7	70.9	66.7	38.6	41.7	24.3	36.1	89.5	22.7	57.1	
rural	28.5	16.3	75.4	NO	31.6	NO	33.6	NO	NO	84.4	59.4	60.8	39.7	35.8	21.8	39.9	42.6	11.9	44.2	
<b>Country region</b>																				
centrale	48.9	27.3	82.7	NO	34.7	NO	35.1	NO	NO	90.2	73.5	75.0	26.6	29.9	20.8	53.6	60.0	14.8	52.7	
kara	28.5	17.8	74.9	NO	30.2	NO	41.0	NO	NO	78.0	56.0	51.9	37.7	28.1	13.3	48.4	63.9	13.6	40.6	
lomé commune	26.1	14.7	93.6	NO	90.8	NO	44.4	NO	NO	95.0	75.8	67.6	37.6	45.2	24.0	31.0	ND	23.9	57.9	
maritime (sans lomé comm)	35.8	22.7	85.0	NO	57.3	NO	34.0	NO	NO	90.4	58.8	62.0	41.9	33.7	24.3	31.9	ND	17.2	50.9	
plateaux	18.5	9.7	84.5	NO	46.9	NO	32.2	NO	NO	87.1	66.9	63.0	45.5	41.5	12.5	41.6	41.9	12.2	45.6	
savanes	15.8	7.5	63.8	NO	16.9	NO	28.2	NO	NO	81.4	56.3	61.2	38.9	55.2	35.0	40.4	48.0	10.7	41.3	




Figure 1 - Coverage levels in the poorest and richest quintiles for selected interventions along the continuum of care.




Figure 2 - Coverage levels in the five wealth quintiles for selected interventions along the continuum of care.




Figure 3 - Co-coverage of health interventions: percentage of children aged 1-4 years according to the number of key child-survival interventions received, by wealth quintile.




Figure 4 - Composite coverage of selected interventions and corresponding coverage gap (how much is needed to reach universal coverage), by wealth quintile.