



Country profiles

The information in the country profiles, customized to showcase the core indicators selected by the Commission on Information and Accountability for Women's and Children's Health, is intended to help policy makers and their partners assess progress and prioritize actions to save women's and children's lives. These country profiles reflect the data that are available for each country. Most of the data for the Commission indicators come from household surveys: mainly the USAID-supported Demographic and Health Surveys and the UNICEF-supported Multiple Indicator Cluster Surveys.^{xxix} More information on the data sources for the Commission indicators is available in the annexes to previous Countdown reports: www.countdown2015mnch.org/reports-and-articles.

New survey data are available for 22 countries since the publication of Countdown's 2012 Report. Coverage estimates that are missing or that are more than five years old indicate an urgent need for concerted action to increase data collection efforts so that updated evidence is available for policy and program development.

This report contains profiles for 75 countries with high burdens of maternal and/or child mortality, including the 49 lowest-income countries covered by the Global Strategy for Women's and Children's Health. Profiles are included for the following countries:

Afghanistan	Gambia	Nigeria
Angola	Ghana	Pakistan
Azerbaijan	Guatemala	Papua New Guinea
Bangladesh	Guinea	Peru
Benin	Guinea-Bissau	Philippines
Bolivia	Haiti	Rwanda
Botswana	India	São Tomé and Príncipe
Brazil	Indonesia	Senegal
Burkina Faso	Iraq	Sierra Leone
Burundi	Kenya	Solomon Islands
Cambodia	Korea, Democratic People's Republic of	Somalia
Cameroon	Kyrgyzstan	South Africa
Central African Republic	Lao People's Democratic Republic	South Sudan
Chad	Lesotho	Sudan
China	Liberia	Swaziland
Comoros	Madagascar	Tajikistan
Congo	Malawi	Tanzania, United Republic of
Congo, Democratic Republic of the	Mali	Togo
Côte d'Ivoire	Mauritania	Turkmenistan
Djibouti	Mexico	Uganda
Egypt	Morocco	Uzbekistan
Equatorial Guinea	Mozambique	Vietnam
Eritrea	Myanmar	Yemen
Ethiopia	Nepal	Zambia
Gabon	Niger	Zimbabwe



Countdown to 2015

Maternal, Newborn & Child Survival

Afghanistan


May 2013

DEMOGRAPHICS

Total population (000)	32,358	(2011)
Total under-five population (000)	5,705	(2011)
Births (000)	1,408	(2011)
Birth registration (%)	37	(2010-2011)
Total under-five deaths (000)	128	(2011)
Neonatal deaths: % of all under-5 deaths	40	(2011)
Neonatal mortality rate (per 1000 live births)	36	(2011)
Infant mortality rate (per 1000 live births)	73	(2011)
Stillbirth rate (per 1000 total births)	29	(2009)
Total maternal deaths	6,400	(2010)
Lifetime risk of maternal death (1 in N)	32	(2010)
Total fertility rate (per woman)	6.2	(2011)
Adolescent birth rate (per 1000 women)	90	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



MICS

NUTRITION

Wasting prevalence (moderate and severe, %) 9 (2004)

Low birthweight incidence (moderate and severe, %) -


Early initiation of breastfeeding (within 1 hr of birth, %) -

Introduction of solid, semi-solid/soft foods (%) 29 (2003)

Vitamin A two dose coverage (%) 100 (2011)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

No Data



Countdown to 2015

Maternal, Newborn & Child Survival

Angola


May 2013

DEMOGRAPHICS

Total population (000)	19,618	(2011)
Total under-five population (000)	3,392	(2011)
Births (000)	803	(2011)
Birth registration (%)	29	(2011)
Total under-five deaths (000)	120	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	43	(2011)
Infant mortality rate (per 1000 live births)	96	(2011)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	3,600	(2010)
Lifetime risk of maternal death (1 in N)	39	(2010)
Total fertility rate (per woman)	5.3	(2011)
Adolescent birth rate (per 1000 women)	165	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



No Data

NUTRITION

Wasting prevalence (moderate and severe, %)

8

(2001)

Low birthweight incidence (moderate and severe, %)

12

(2001)

Early initiation of breastfeeding (within 1 hr of birth, %)

-

Introduction of solid, semi-solid/soft foods (%)

-


Vitamin A two dose coverage (%)

55

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Azerbaijan


May 2013

DEMOGRAPHICS

Total population (000)	9,306	(2011)
Total under-five population (000)	846	(2011)
Births (000)	184	(2011)
Birth registration (%)	94	(2006)
Total under-five deaths (000)	8	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	19	(2011)
Infant mortality rate (per 1000 live births)	39	(2011)
Stillbirth rate (per 1000 total births)	12	(2009)
Total maternal deaths	79	(2010)
Lifetime risk of maternal death (1 in N)	1,000	(2010)
Total fertility rate (per woman)	2.2	(2011)
Adolescent birth rate (per 1000 women)	41	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS

NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2006)

Low birthweight incidence (moderate and severe, %) 10 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 32 (2006)

Introduction of solid, semi-solid/soft foods (%) 83 (2006)

Vitamin A two dose coverage (%) 89 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Bangladesh


May 2013

DEMOGRAPHICS

Total population (000)	150,494	(2011)
Total under-five population (000)	14,427	(2011)
Births (000)	3,016	(2011)
Birth registration (%)	10	(2006)
Total under-five deaths (000)	134	(2011)
Neonatal deaths: % of all under-5 deaths	60	(2011)
Neonatal mortality rate (per 1000 live births)	26	(2011)
Infant mortality rate (per 1000 live births)	37	(2011)
Stillbirth rate (per 1000 total births)	36	(2009)
Total maternal deaths	7,200	(2010)
Lifetime risk of maternal death (1 in N)	170	(2010)
Total fertility rate (per woman)	2.2	(2011)
Adolescent birth rate (per 1000 women)	133	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 16 (2011)
Low birthweight incidence (moderate and severe, %) 22 (2006)

Early initiation of breastfeeding (within 1 hr of birth, %) 36 (2006)
Introduction of solid, semi-solid/soft foods (%) 71 (2007)
Vitamin A two dose coverage (%) 94 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Benin


May 2013

DEMOGRAPHICS

Total population (000)	9,100	(2011)
Total under-five population (000)	1,547	(2011)
Births (000)	356	(2011)
Birth registration (%)	60	(2011)
Total under-five deaths (000)	36	(2011)
Neonatal deaths: % of all under-5 deaths	30	(2011)
Neonatal mortality rate (per 1000 live births)	31	(2011)
Infant mortality rate (per 1000 live births)	68	(2011)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	53	(2010)
Total fertility rate (per woman)	5.2	(2011)
Adolescent birth rate (per 1000 women)	114	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%



DHS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

8

(2006)

Low birthweight incidence (moderate and severe, %)

15

(2006)

Early initiation of breastfeeding (within 1 hr of birth, %)

32

(2008)

Introduction of solid, semi-solid/soft foods (%)

76

(2008)


Vitamin A two dose coverage (%)

98

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Bolivia


May 2013

DEMOGRAPHICS

Total population (000)	10,088	(2011)
Total under-five population (000)	1,229	(2011)
Births (000)	264	(2011)
Birth registration (%)	76	(2008)
Total under-five deaths (000)	13	(2011)
Neonatal deaths: % of all under-5 deaths	45	(2011)
Neonatal mortality rate (per 1000 live births)	22	(2011)
Infant mortality rate (per 1000 live births)	39	(2011)
Stillbirth rate (per 1000 total births)	17	(2009)
Total maternal deaths	510	(2010)
Lifetime risk of maternal death (1 in N)	140	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	89	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

1

(2008)

Low birthweight incidence (moderate and severe, %)

6

(2008)

Early initiation of breastfeeding (within 1 hr of birth, %)

64

(2008)

Introduction of solid, semi-solid/soft foods (%)

83

(2008)


Vitamin A two dose coverage (%)

21

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Botswana


May 2013

DEMOGRAPHICS

Total population (000)	2,031	(2011)
Total under-five population (000)	229	(2011)
Births (000)	47	(2011)
Birth registration (%)	72	(2007)
Total under-five deaths (000)	1	(2011)
Neonatal deaths: % of all under-5 deaths	43	(2011)
Neonatal mortality rate (per 1000 live births)	11	(2011)
Infant mortality rate (per 1000 live births)	20	(2011)
Stillbirth rate (per 1000 total births)	16	(2009)
Total maternal deaths	75	(2010)
Lifetime risk of maternal death (1 in N)	220	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	51	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS

NUTRITION

Wasting prevalence (moderate and severe, %)

7

(2007)

Low birthweight incidence (moderate and severe, %)

13

(2007)

Early initiation of breastfeeding (within 1 hr of birth, %)

40

(2007)

Introduction of solid, semi-solid/soft foods (%)

46

(2007)


Vitamin A two dose coverage (%)

75

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Brazil


May 2013

DEMOGRAPHICS

Total population (000)	196,655	(2011)
Total under-five population (000)	14,647	(2011)
Births (000)	2,996	(2011)
Birth registration (%)	91	(2008)
Total under-five deaths (000)	44	(2011)
Neonatal deaths: % of all under-5 deaths	66	(2011)
Neonatal mortality rate (per 1000 live births)	10	(2011)
Infant mortality rate (per 1000 live births)	14	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	1,700	(2010)
Lifetime risk of maternal death (1 in N)	910	(2010)
Total fertility rate (per woman)	1.8	(2011)
Adolescent birth rate (per 1000 women)	71	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 1996


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 2 (2006)

Low birthweight incidence (moderate and severe, %) 8 (2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 43 (2006)

Introduction of solid, semi-solid/soft foods (%) 70 (2008)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Burkina Faso


May 2013

DEMOGRAPHICS

Total population (000)	16,968	(2011)
Total under-five population (000)	3,051	(2011)
Births (000)	730	(2011)
Birth registration (%)	77	(2010)
Total under-five deaths (000)	101	(2011)
Neonatal deaths: % of all under-5 deaths	25	(2011)
Neonatal mortality rate (per 1000 live births)	34	(2011)
Infant mortality rate (per 1000 live births)	82	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	2,100	(2010)
Lifetime risk of maternal death (1 in N)	55	(2010)
Total fertility rate (per woman)	5.8	(2011)
Adolescent birth rate (per 1000 women)	130	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT


Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.


CHILD HEALTH

Immunization

Percent of children immunized against measles

Percent of children immunized with 3 doses DTP

Percent of children immunized with 3 doses Hib




Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider

Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

11

(2009)

Low birthweight incidence (moderate and severe, %)

16

(2006)

Early initiation of breastfeeding (within 1 hr of birth, %)

20

(2006)

Introduction of solid, semi-solid/soft foods (%)

61

(2010-2011)

Vitamin A two dose coverage (%)


87

(2011)

Underweight and stunting prevalence

Percent children <5 years who are underweight


Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Burundi


May 2013

DEMOGRAPHICS

Total population (000)	8,575	(2011)
Total under-five population (000)	1,218	(2011)
Births (000)	288	(2011)
Birth registration (%)	75	(2010)
Total under-five deaths (000)	39	(2011)
Neonatal deaths: % of all under-5 deaths	32	(2011)
Neonatal mortality rate (per 1000 live births)	43	(2011)
Infant mortality rate (per 1000 live births)	86	(2011)
Stillbirth rate (per 1000 total births)	28	(2009)
Total maternal deaths	2,200	(2010)
Lifetime risk of maternal death (1 in N)	31	(2010)
Total fertility rate (per woman)	4.2	(2011)
Adolescent birth rate (per 1000 women)	65	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS, MICS, Other NS


NUTRITION

Wasting prevalence (moderate and severe, %)	6	(2010)
Low birthweight incidence (moderate and severe, %)	11	(2005)

Early initiation of breastfeeding (within 1 hr of birth, %)	-	(2010)
Introduction of solid, semi-solid/soft foods (%)	70	(2010)
Vitamin A two dose coverage (%)	83	(2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Cambodia


May 2013

DEMOGRAPHICS

Total population (000)	14,305	(2011)
Total under-five population (000)	1,507	(2011)
Births (000)	317	(2011)
Birth registration (%)	62	(2010)
Total under-five deaths (000)	13	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	19	(2011)
Infant mortality rate (per 1000 live births)	36	(2011)
Stillbirth rate (per 1000 total births)	18	(2009)
Total maternal deaths	790	(2010)
Lifetime risk of maternal death (1 in N)	150	(2010)
Total fertility rate (per woman)	2.5	(2011)
Adolescent birth rate (per 1000 women)	48	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 11 (2010)

Low birthweight incidence (moderate and severe, %) 11 (2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 65 (2010)

Introduction of solid, semi-solid/soft foods (%) 82 (2010)

Vitamin A two dose coverage (%) 92 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Cameroon


May 2013

DEMOGRAPHICS

Total population (000)	20,030	(2011)
Total under-five population (000)	3,104	(2011)
Births (000)	716	(2011)
Birth registration (%)	70	(2006)
Total under-five deaths (000)	88	(2011)
Neonatal deaths: % of all under-5 deaths	27	(2011)
Neonatal mortality rate (per 1000 live births)	33	(2011)
Infant mortality rate (per 1000 live births)	79	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	4,900	(2010)
Lifetime risk of maternal death (1 in N)	31	(2010)
Total fertility rate (per woman)	4.4	(2011)
Adolescent birth rate (per 1000 women)	127	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2011


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 6 (2011)
Low birthweight incidence (moderate and severe, %) 11 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Central African Republic


May 2013

DEMOGRAPHICS

Total population (000)	4,487	(2011)
Total under-five population (000)	659	(2011)
Births (000)	156	(2011)
Birth registration (%)	61	(2010)
Total under-five deaths (000)	25	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	46	(2011)
Infant mortality rate (per 1000 live births)	108	(2011)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	1,400	(2010)
Lifetime risk of maternal death (1 in N)	26	(2010)
Total fertility rate (per woman)	4.5	(2011)
Adolescent birth rate (per 1000 women)	133	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2010)

Low birthweight incidence (moderate and severe, %) 14 (2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 43 (2010)

Introduction of solid, semi-solid/soft foods (%) 56 (2010)

Vitamin A two dose coverage (%) 0 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Chad


May 2013

DEMOGRAPHICS

Total population (000)	11,525	(2011)
Total under-five population (000)	2,049	(2011)
Births (000)	511	(2011)
Birth registration (%)	16	(2011)
Total under-five deaths (000)	79	(2011)
Neonatal deaths: % of all under-5 deaths	27	(2011)
Neonatal mortality rate (per 1000 live births)	42	(2011)
Infant mortality rate (per 1000 live births)	97	(2011)
Stillbirth rate (per 1000 total births)	29	(2009)
Total maternal deaths	5,300	(2010)
Lifetime risk of maternal death (1 in N)	15	(2010)
Total fertility rate (per woman)	5.9	(2011)
Adolescent birth rate (per 1000 women)	193	(2002)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 2004


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 16 (2010)
Low birthweight incidence (moderate and severe, %) 20 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

China


May 2013

DEMOGRAPHICS

Total population (000)	1,347,565	(2011)
Total under-five population (000)	82,231	(2011)
Births (000)	16,364	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	249	(2011)
Neonatal deaths: % of all under-5 deaths	58	(2011)
Neonatal mortality rate (per 1000 live births)	9	(2011)
Infant mortality rate (per 1000 live births)	13	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	6,000	(2010)
Lifetime risk of maternal death (1 in N)	1,700	(2010)
Total fertility rate (per woman)	1.6	(2011)
Adolescent birth rate (per 1000 women)	6	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: MOH

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT


Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics

No Data

NUTRITION

Wasting prevalence (moderate and severe, %)

3 (2002)

Low birthweight incidence (moderate and severe, %)

3 (2008)

Early initiation of breastfeeding (within 1 hr of birth, %)

41 (2008)

Introduction of solid, semi-solid/soft foods (%)


43 (2008)

Vitamin A two dose coverage (%)

-

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Comoros


May 2013

DEMOGRAPHICS

Total population (000)	754	(2011)
Total under-five population (000)	124	(2011)
Births (000)	28	(2011)
Birth registration (%)	83	(2011)
Total under-five deaths (000)	2	(2011)
Neonatal deaths: % of all under-5 deaths	41	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	59	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	79	(2010)
Lifetime risk of maternal death (1 in N)	67	(2010)
Total fertility rate (per woman)	4.9	(2011)
Adolescent birth rate (per 1000 women)	95	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

MICS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 1996


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



DHS

MICS

NUTRITION

Wasting prevalence (moderate and severe, %) - -

Low birthweight incidence (moderate and severe, %) 25 (2000)


Early initiation of breastfeeding (within 1 hr of birth, %) 25 (1996)

Introduction of solid, semi-solid/soft foods (%) 34 (2000)

Vitamin A two dose coverage (%) 18 (2010)


Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



DHS

MICS



Countdown to 2015

Maternal, Newborn & Child Survival

Congo


May 2013

DEMOGRAPHICS

Total population (000)	4,140	(2011)
Total under-five population (000)	636	(2011)
Births (000)	145	(2011)
Birth registration (%)	81	(2009)
Total under-five deaths (000)	14	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	64	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	800	(2010)
Lifetime risk of maternal death (1 in N)	39	(2010)
Total fertility rate (per woman)	4.5	(2011)
Adolescent birth rate (per 1000 women)	132	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



pDHS

NUTRITION

Wasting prevalence (moderate and severe, %) 8 (2005)

Low birthweight incidence (moderate and severe, %) 13 (2005)


Early initiation of breastfeeding (within 1 hr of birth, %) 39 (2005)

Introduction of solid, semi-solid/soft foods (%) 78 (2005)

Vitamin A two dose coverage (%) 84 (2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted




Other NS

DHS

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



DHS

Note: Based on 2006 WHO reference population

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Congo, Democratic Republic of the


May 2013

DEMOGRAPHICS

Total population (000)	67,758	(2011)
Total under-five population (000)	12,046	(2011)
Births (000)	2,912	(2011)
Birth registration (%)	28	(2010)
Total under-five deaths (000)	465	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	47	(2011)
Infant mortality rate (per 1000 live births)	111	(2011)
Stillbirth rate (per 1000 total births)	29	(2009)
Total maternal deaths	15,000	(2010)
Lifetime risk of maternal death (1 in N)	30	(2010)
Total fertility rate (per woman)	5.7	(2011)
Adolescent birth rate (per 1000 women)	135	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2010


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 9 (2010)
Low birthweight incidence (moderate and severe, %) 10 (2010)

Early initiation of breastfeeding (within 1 hr of birth, %) 43 (2010)
Introduction of solid, semi-solid/soft foods (%) 52 (2010)
Vitamin A two dose coverage (%) 98 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Côte d'Ivoire


May 2013

DEMOGRAPHICS

Total population (000)	20,153	(2011)
Total under-five population (000)	2,996	(2011)
Births (000)	679	(2011)
Birth registration (%)	55	(2009)
Total under-five deaths (000)	75	(2011)
Neonatal deaths: % of all under-5 deaths	37	(2011)
Neonatal mortality rate (per 1000 live births)	41	(2011)
Infant mortality rate (per 1000 live births)	81	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	2,700	(2010)
Lifetime risk of maternal death (1 in N)	53	(2010)
Total fertility rate (per woman)	4.3	(2011)
Adolescent birth rate (per 1000 women)	111	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2011)

Low birthweight incidence (moderate and severe, %) 17 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 25 (2006)

Introduction of solid, semi-solid/soft foods (%) 51 (2006)

Vitamin A two dose coverage (%) 100 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Djibouti


May 2013

DEMOGRAPHICS

Total population (000)	906	(2011)
Total under-five population (000)	115	(2011)
Births (000)	26	(2011)
Birth registration (%)	89	(2006)
Total under-five deaths (000)	2	(2011)
Neonatal deaths: % of all under-5 deaths	38	(2011)
Neonatal mortality rate (per 1000 live births)	33	(2011)
Infant mortality rate (per 1000 live births)	72	(2011)
Stillbirth rate (per 1000 total births)	34	(2009)
Total maternal deaths	51	(2010)
Lifetime risk of maternal death (1 in N)	140	(2010)
Total fertility rate (per woman)	3.7	(2011)
Adolescent birth rate (per 1000 women)	27	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2010)
Low birthweight incidence (moderate and severe, %) 10 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Egypt


May 2013

DEMOGRAPHICS

Total population (000)	82,537	(2011)
Total under-five population (000)	9,093	(2011)
Births (000)	1,886	(2011)
Birth registration (%)	99	(2005)
Total under-five deaths (000)	40	(2011)
Neonatal deaths: % of all under-5 deaths	35	(2011)
Neonatal mortality rate (per 1000 live births)	7	(2011)
Infant mortality rate (per 1000 live births)	18	(2011)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	490	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	50	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2008)

Low birthweight incidence (moderate and severe, %) 13 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 56 (2008)

Introduction of solid, semi-solid/soft foods (%) 70 (2008)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Equatorial Guinea


May 2013

DEMOGRAPHICS

Total population (000)	720	(2011)
Total under-five population (000)	111	(2011)
Births (000)	26	(2011)
Birth registration (%)	32	(2011)
Total under-five deaths (000)	3	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	37	(2011)
Infant mortality rate (per 1000 live births)	80	(2011)
Stillbirth rate (per 1000 total births)	17	(2009)
Total maternal deaths	61	(2010)
Lifetime risk of maternal death (1 in N)	88	(2010)
Total fertility rate (per woman)	5.1	(2011)
Adolescent birth rate (per 1000 women)	128	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics

No Data


NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2004)
Low birthweight incidence (moderate and severe, %)	13	(2000)

Early initiation of breastfeeding (within 1 hr of birth, %)	-	-
Introduction of solid, semi-solid/soft foods (%)	-	-
Vitamin A two dose coverage (%)	0	(2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Eritrea


May 2013

DEMOGRAPHICS

Total population (000)	5,415	(2011)
Total under-five population (000)	880	(2011)
Births (000)	193	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	13	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	22	(2011)
Infant mortality rate (per 1000 live births)	46	(2011)
Stillbirth rate (per 1000 total births)	21	(2009)
Total maternal deaths	460	(2010)
Lifetime risk of maternal death (1 in N)	86	(2010)
Total fertility rate (per woman)	4.4	(2011)
Adolescent birth rate (per 1000 women)	85	(1999)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2002)

Low birthweight incidence (moderate and severe, %) 14 (2002)


Early initiation of breastfeeding (within 1 hr of birth, %) 78 (2002)

Introduction of solid, semi-solid/soft foods (%) 43 (2002)

Vitamin A two dose coverage (%) 46 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: DHS



Countdown to 2015

Maternal, Newborn & Child Survival

Ethiopia


May 2013

DEMOGRAPHICS

Total population (000)	84,734	(2011)
Total under-five population (000)	11,918	(2011)
Births (000)	2,613	(2011)
Birth registration (%)	7	(2005)
Total under-five deaths (000)	194	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	31	(2011)
Infant mortality rate (per 1000 live births)	52	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	9,000	(2010)
Lifetime risk of maternal death (1 in N)	67	(2010)
Total fertility rate (per woman)	4.0	(2011)
Adolescent birth rate (per 1000 women)	79	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2011

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2011)
Low birthweight incidence (moderate and severe, %) 20 (2005)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Gabon


May 2013

DEMOGRAPHICS

Total population (000)	1,534	(2011)
Total under-five population (000)	188	(2011)
Births (000)	42	(2011)
Birth registration (%)	89	(2008)
Total under-five deaths (000)	3	(2011)
Neonatal deaths: % of all under-5 deaths	39	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	49	(2011)
Stillbirth rate (per 1000 total births)	17	(2009)
Total maternal deaths	94	(2010)
Lifetime risk of maternal death (1 in N)	130	(2010)
Total fertility rate (per woman)	3.2	(2011)
Adolescent birth rate (per 1000 women)	144	(1998)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2000

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2000)

Low birthweight incidence (moderate and severe, %) 14 (2000)


Early initiation of breastfeeding (within 1 hr of birth, %) 71 (2000)

Introduction of solid, semi-solid/soft foods (%) 62 (2000)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Gambia


May 2013

DEMOGRAPHICS

Total population (000)	1,776	(2011)
Total under-five population (000)	292	(2011)
Births (000)	67	(2011)
Birth registration (%)	53	(2010)
Total under-five deaths (000)	6	(2011)
Neonatal deaths: % of all under-5 deaths	36	(2011)
Neonatal mortality rate (per 1000 live births)	34	(2011)
Infant mortality rate (per 1000 live births)	58	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	230	(2010)
Lifetime risk of maternal death (1 in N)	56	(2010)
Total fertility rate (per woman)	4.8	(2011)
Adolescent birth rate (per 1000 women)	104	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2005-2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2010)
Low birthweight incidence (moderate and severe, %) 10 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Ghana


May 2013

DEMOGRAPHICS

Total population (000)	24,966	(2011)
Total under-five population (000)	3,592	(2011)
Births (000)	776	(2011)
Birth registration (%)	63	(2011)
Total under-five deaths (000)	60	(2011)
Neonatal deaths: % of all under-5 deaths	38	(2011)
Neonatal mortality rate (per 1000 live births)	30	(2011)
Infant mortality rate (per 1000 live births)	52	(2011)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	2,700	(2010)
Lifetime risk of maternal death (1 in N)	68	(2010)
Total fertility rate (per woman)	4.1	(2011)
Adolescent birth rate (per 1000 women)	70	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 9 (2008)

Low birthweight incidence (moderate and severe, %) 13 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 52 (2008)

Introduction of solid, semi-solid/soft foods (%) 76 (2008)

Vitamin A two dose coverage (%) 93 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Guatemala


May 2013

DEMOGRAPHICS

Total population (000)	14,757	(2011)
Total under-five population (000)	2,192	(2011)
Births (000)	473	(2011)
Birth registration (%)	97	(2008-2009)
Total under-five deaths (000)	14	(2011)
Neonatal deaths: % of all under-5 deaths	49	(2011)
Neonatal mortality rate (per 1000 live births)	15	(2011)
Infant mortality rate (per 1000 live births)	24	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	550	(2010)
Lifetime risk of maternal death (1 in N)	190	(2010)
Total fertility rate (per woman)	3.9	(2011)
Adolescent birth rate (per 1000 women)	92	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



DHS 1998-1999


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 1 (2008-2009)

Low birthweight incidence (moderate and severe, %) 11 (2008-2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 56 (2008-2009)

Introduction of solid, semi-solid/soft foods (%) 71 (2008-2009)

Vitamin A two dose coverage (%) 28 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Guinea


May 2013

DEMOGRAPHICS

Total population (000)	10,222	(2011)
Total under-five population (000)	1,696	(2011)
Births (000)	394	(2011)
Birth registration (%)	43	(2005)
Total under-five deaths (000)	48	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	39	(2011)
Infant mortality rate (per 1000 live births)	79	(2011)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	2,400	(2010)
Lifetime risk of maternal death (1 in N)	30	(2010)
Total fertility rate (per woman)	5.2	(2011)
Adolescent birth rate (per 1000 women)	153	(2002)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2005


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 8 (2008)
Low birthweight incidence (moderate and severe, %) 12 (2005)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Guinea-Bissau


May 2013

DEMOGRAPHICS

Total population (000)	1,547	(2011)
Total under-five population (000)	245	(2011)
Births (000)	59	(2011)
Birth registration (%)	24	(2010)
Total under-five deaths (000)	9	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	44	(2011)
Infant mortality rate (per 1000 live births)	98	(2011)
Stillbirth rate (per 1000 total births)	30	(2009)
Total maternal deaths	460	(2010)
Lifetime risk of maternal death (1 in N)	25	(2010)
Total fertility rate (per woman)	5.0	(2011)
Adolescent birth rate (per 1000 women)	137	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 6 (2010)
Low birthweight incidence (moderate and severe, %) 11 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Haiti


May 2013

DEMOGRAPHICS

Total population (000)	10,124	(2011)
Total under-five population (000)	1,246	(2011)
Births (000)	266	(2011)
Birth registration (%)	81	(2005-2006)
Total under-five deaths (000)	19	(2011)
Neonatal deaths: % of all under-5 deaths	36	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	940	(2010)
Lifetime risk of maternal death (1 in N)	83	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	69	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2005-2006)

Low birthweight incidence (moderate and severe, %) 25 (2005-2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 44 (2005-2006)

Introduction of solid, semi-solid/soft foods (%) 90 (2005-2006)

Vitamin A two dose coverage (%) 36 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

India


May 2013

DEMOGRAPHICS

Total population (000)	1,241,492	(2011)
Total under-five population (000)	128,589	(2011)
Births (000)	27,098	(2011)
Birth registration (%)	41	(2005-2006)
Total under-five deaths (000)	1,655	(2011)
Neonatal deaths: % of all under-5 deaths	53	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	47	(2011)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	56,000	(2010)
Lifetime risk of maternal death (1 in N)	170	(2010)
Total fertility rate (per woman)	2.6	(2011)
Adolescent birth rate (per 1000 women)	39	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




NFHS 2005-2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 20 (2005-2006)
Low birthweight incidence (moderate and severe, %) 28 (2005-2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Indonesia


May 2013

DEMOGRAPHICS

Total population (000)	242,326	(2011)
Total under-five population (000)	21,199	(2011)
Births (000)	4,331	(2011)
Birth registration (%)	53	(2007)
Total under-five deaths (000)	134	(2011)
Neonatal deaths: % of all under-5 deaths	49	(2011)
Neonatal mortality rate (per 1000 live births)	15	(2011)
Infant mortality rate (per 1000 live births)	25	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	9,600	(2010)
Lifetime risk of maternal death (1 in N)	210	(2010)
Total fertility rate (per woman)	2.1	(2011)
Adolescent birth rate (per 1000 women)	52	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 13 (2010)

Low birthweight incidence (moderate and severe, %) 9 (2007)


Early initiation of breastfeeding (within 1 hr of birth, %) 29 (2010)

Introduction of solid, semi-solid/soft foods (%) 85 (2007)

Vitamin A two dose coverage (%) 76 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Iraq


May 2013

DEMOGRAPHICS

Total population (000)	32,665	(2011)
Total under-five population (000)	5,301	(2011)
Births (000)	1,144	(2011)
Birth registration (%)	95	(2006)
Total under-five deaths (000)	42	(2011)
Neonatal deaths: % of all under-5 deaths	54	(2011)
Neonatal mortality rate (per 1000 live births)	20	(2011)
Infant mortality rate (per 1000 live births)	31	(2011)
Stillbirth rate (per 1000 total births)	9	(2009)
Total maternal deaths	710	(2010)
Lifetime risk of maternal death (1 in N)	310	(2010)
Total fertility rate (per woman)	4.6	(2011)
Adolescent birth rate (per 1000 women)	68	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 6 (2006)
Low birthweight incidence (moderate and severe, %) 15 (2006)

Early initiation of breastfeeding (within 1 hr of birth, %) 31 (2006)
Introduction of solid, semi-solid/soft foods (%) 62 (2006)
Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Kenya


May 2013

DEMOGRAPHICS

Total population (000)	41,610	(2011)
Total under-five population (000)	6,807	(2011)
Births (000)	1,560	(2011)
Birth registration (%)	60	(2008-2009)
Total under-five deaths (000)	107	(2011)
Neonatal deaths: % of all under-5 deaths	39	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	48	(2011)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	5,500	(2010)
Lifetime risk of maternal death (1 in N)	55	(2010)
Total fertility rate (per woman)	4.7	(2011)
Adolescent birth rate (per 1000 women)	106	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2008-2009)

Low birthweight incidence (moderate and severe, %) 8 (2008-2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 58 (2008-2009)

Introduction of solid, semi-solid/soft foods (%) 85 (2008-2009)

Vitamin A two dose coverage (%) 62 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Korea, Democratic People's Republic of


May 2013

DEMOGRAPHICS

Total population (000)	24,451	(2011)
Total under-five population (000)	1,706	(2011)
Births (000)	348	(2011)
Birth registration (%)	100	(2009)
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	52	(2011)
Neonatal mortality rate (per 1000 live births)	18	(2011)
Infant mortality rate (per 1000 live births)	26	(2011)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	280	(2010)
Lifetime risk of maternal death (1 in N)	670	(2010)
Total fertility rate (per woman)	2.0	(2011)
Adolescent birth rate (per 1000 women)	1	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2009)

Low birthweight incidence (moderate and severe, %) 6 (2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 18 (2009)

Introduction of solid, semi-solid/soft foods (%) 31 (2004)

Vitamin A two dose coverage (%) 100 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Kyrgyzstan


May 2013

DEMOGRAPHICS

Total population (000)	5,393	(2011)
Total under-five population (000)	626	(2011)
Births (000)	131	(2011)
Birth registration (%)	94	(2009)
Total under-five deaths (000)	4	(2011)
Neonatal deaths: % of all under-5 deaths	48	(2011)
Neonatal mortality rate (per 1000 live births)	16	(2011)
Infant mortality rate (per 1000 live births)	27	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	91	(2010)
Lifetime risk of maternal death (1 in N)	480	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	31	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 3 (2006)

Low birthweight incidence (moderate and severe, %) 5 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 65 (2006)

Introduction of solid, semi-solid/soft foods (%) 60 (2006)

Vitamin A two dose coverage (%) 97 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Lao People's Democratic Republic


May 2013

DEMOGRAPHICS

Total population (000)	6,288	(2011)
Total under-five population (000)	682	(2011)
Births (000)	140	(2011)
Birth registration (%)	72	(2006)
Total under-five deaths (000)	6	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	18	(2011)
Infant mortality rate (per 1000 live births)	34	(2011)
Stillbirth rate (per 1000 total births)	14	(2009)
Total maternal deaths	670	(2010)
Lifetime risk of maternal death (1 in N)	74	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	110	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2006)
Low birthweight incidence (moderate and severe, %) 11 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Lesotho


May 2013

DEMOGRAPHICS

Total population (000)	2,194	(2011)
Total under-five population (000)	276	(2011)
Births (000)	60	(2011)
Birth registration (%)	45	(2009)
Total under-five deaths (000)	5	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	39	(2011)
Infant mortality rate (per 1000 live births)	63	(2011)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	370	(2010)
Lifetime risk of maternal death (1 in N)	53	(2010)
Total fertility rate (per woman)	3.1	(2011)
Adolescent birth rate (per 1000 women)	92	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2009


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS, DHS

NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2009)

Low birthweight incidence (moderate and severe, %) 11 (2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 53 (2009)

Introduction of solid, semi-solid/soft foods (%) 68 (2009)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: MICS, DHS



Countdown to 2015

Maternal, Newborn & Child Survival

Liberia


May 2013

DEMOGRAPHICS

Total population (000)	4,129	(2011)
Total under-five population (000)	696	(2011)
Births (000)	157	(2011)
Birth registration (%)	4	(2007)
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	36	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	58	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	24	(2010)
Total fertility rate (per woman)	5.2	(2011)
Adolescent birth rate (per 1000 women)	177	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2010)
Low birthweight incidence (moderate and severe, %)	14	(2007)

Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2010)
Introduction of solid, semi-solid/soft foods (%)	51	(2010)
Vitamin A two dose coverage (%)	96	(2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Madagascar


May 2013

DEMOGRAPHICS

Total population (000)	21,315	(2011)
Total under-five population (000)	3,379	(2011)
Births (000)	747	(2011)
Birth registration (%)	80	(2008-2009)
Total under-five deaths (000)	45	(2011)
Neonatal deaths: % of all under-5 deaths	38	(2011)
Neonatal mortality rate (per 1000 live births)	23	(2011)
Infant mortality rate (per 1000 live births)	43	(2011)
Stillbirth rate (per 1000 total births)	21	(2009)
Total maternal deaths	1,800	(2010)
Lifetime risk of maternal death (1 in N)	81	(2010)
Total fertility rate (per woman)	4.6	(2011)
Adolescent birth rate (per 1000 women)	147	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2003-2004)
Low birthweight incidence (moderate and severe, %) 16 (2008-2009)

Early initiation of breastfeeding (within 1 hr of birth, %) 72 (2008-2009)
Introduction of solid, semi-solid/soft foods (%) 86 (2008-2009)
Vitamin A two dose coverage (%) 91 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Malawi


May 2013

DEMOGRAPHICS

Total population (000)	15,381	(2011)
Total under-five population (000)	2,832	(2011)
Births (000)	686	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	52	(2011)
Neonatal deaths: % of all under-5 deaths	35	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	3,000	(2010)
Lifetime risk of maternal death (1 in N)	36	(2010)
Total fertility rate (per woman)	6.0	(2011)
Adolescent birth rate (per 1000 women)	157	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2010)

Low birthweight incidence (moderate and severe, %) 13 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 58 (2006)

Introduction of solid, semi-solid/soft foods (%) 86 (2010)

Vitamin A two dose coverage (%) 96 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mali


May 2013

DEMOGRAPHICS

Total population (000)	15,840	(2011)
Total under-five population (000)	2,998	(2011)
Births (000)	728	(2011)
Birth registration (%)	81	(2010)
Total under-five deaths (000)	121	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	49	(2011)
Infant mortality rate (per 1000 live births)	98	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	3,800	(2010)
Lifetime risk of maternal death (1 in N)	28	(2010)
Total fertility rate (per woman)	6.2	(2011)
Adolescent birth rate (per 1000 women)	190	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%



DHS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

15

(2006)

Low birthweight incidence (moderate and severe, %)

19

(2006)

Early initiation of breastfeeding (within 1 hr of birth, %)

46

(2006)

Introduction of solid, semi-solid/soft foods (%)

25

(2006)


Vitamin A two dose coverage (%)

96

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Mauritania


May 2013

DEMOGRAPHICS

Total population (000)	3,542	(2011)
Total under-five population (000)	522	(2011)
Births (000)	118	(2011)
Birth registration (%)	56	(2007)
Total under-five deaths (000)	13	(2011)
Neonatal deaths: % of all under-5 deaths	37	(2011)
Neonatal mortality rate (per 1000 live births)	40	(2011)
Infant mortality rate (per 1000 live births)	76	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	590	(2010)
Lifetime risk of maternal death (1 in N)	44	(2010)
Total fertility rate (per woman)	4.5	(2011)
Adolescent birth rate (per 1000 women)	88	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 12 (2010)
Low birthweight incidence (moderate and severe, %) 34 (2007)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Mexico


May 2013

DEMOGRAPHICS

Total population (000)	114,793	(2011)
Total under-five population (000)	10,938	(2011)
Births (000)	2,195	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	34	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	7	(2011)
Infant mortality rate (per 1000 live births)	13	(2011)
Stillbirth rate (per 1000 total births)	5	(2009)
Total maternal deaths	1,100	(2010)
Lifetime risk of maternal death (1 in N)	790	(2010)
Total fertility rate (per woman)	2.3	(2011)
Adolescent birth rate (per 1000 women)	87	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: UNICEF/UNAIDS/WHO

NUTRITION

Wasting prevalence (moderate and severe, %) - -

Low birthweight incidence (moderate and severe, %) 7 (2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 18 (2009)

Introduction of solid, semi-solid/soft foods (%) 27 (2009)

Vitamin A two dose coverage (%) - -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Morocco


May 2013

DEMOGRAPHICS

Total population (000)	32,273	(2011)
Total under-five population (000)	3,051	(2011)
Births (000)	620	(2011)
Birth registration (%)	85	(2000)
Total under-five deaths (000)	21	(2011)
Neonatal deaths: % of all under-5 deaths	56	(2011)
Neonatal mortality rate (per 1000 live births)	19	(2011)
Infant mortality rate (per 1000 live births)	28	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	650	(2010)
Lifetime risk of maternal death (1 in N)	400	(2010)
Total fertility rate (per woman)	2.2	(2011)
Adolescent birth rate (per 1000 women)	18	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2003-2004


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS, Other NS

NUTRITION

Wasting prevalence (moderate and severe, %)

2 (2010-2011)

Low birthweight incidence (moderate and severe, %)

15 (2003-2004)

Early initiation of breastfeeding (within 1 hr of birth, %)

52 (2003-2004)

Introduction of solid, semi-solid/soft foods (%)


66 (2003-2004)

Vitamin A two dose coverage (%)

-

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Mozambique


May 2013

DEMOGRAPHICS

Total population (000)	23,930	(2011)
Total under-five population (000)	3,875	(2011)
Births (000)	889	(2011)
Birth registration (%)	31	(2008)
Total under-five deaths (000)	86	(2011)
Neonatal deaths: % of all under-5 deaths	35	(2011)
Neonatal mortality rate (per 1000 live births)	34	(2011)
Infant mortality rate (per 1000 live births)	72	(2011)
Stillbirth rate (per 1000 total births)	28	(2009)
Total maternal deaths	4,300	(2010)
Lifetime risk of maternal death (1 in N)	43	(2010)
Total fertility rate (per woman)	4.8	(2011)
Adolescent birth rate (per 1000 women)	193	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 6 (2011)
Low birthweight incidence (moderate and severe, %) 16 (2008)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Myanmar


May 2013

DEMOGRAPHICS

Total population (000)	48,337	(2011)
Total under-five population (000)	3,985	(2011)
Births (000)	824	(2011)
Birth registration (%)	72	(2009-2010)
Total under-five deaths (000)	53	(2011)
Neonatal deaths: % of all under-5 deaths	47	(2011)
Neonatal mortality rate (per 1000 live births)	30	(2011)
Infant mortality rate (per 1000 live births)	48	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	1,600	(2010)
Lifetime risk of maternal death (1 in N)	250	(2010)
Total fertility rate (per woman)	2.0	(2011)
Adolescent birth rate (per 1000 women)	17	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS

NUTRITION

Wasting prevalence (moderate and severe, %) 8 (2009-2010)

Low birthweight incidence (moderate and severe, %) 9 (2009-2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 76 (2009-2010)

Introduction of solid, semi-solid/soft foods (%) 81 (2009-2010)

Vitamin A two dose coverage (%) 96 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: MICS



Countdown to 2015

Maternal, Newborn & Child Survival

Nepal


May 2013

DEMOGRAPHICS

Total population (000)	30,486	(2011)
Total under-five population (000)	3,450	(2011)
Births (000)	722	(2011)
Birth registration (%)	42	(2011)
Total under-five deaths (000)	34	(2011)
Neonatal deaths: % of all under-5 deaths	58	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	39	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	190	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	81	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2011

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 11 (2011)
Low birthweight incidence (moderate and severe, %) 18 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Niger


May 2013

DEMOGRAPHICS

Total population (000)	16,069	(2011)
Total under-five population (000)	3,200	(2011)
Births (000)	777	(2011)
Birth registration (%)	32	(2006)
Total under-five deaths (000)	89	(2011)
Neonatal deaths: % of all under-5 deaths	28	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	66	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	4,500	(2010)
Lifetime risk of maternal death (1 in N)	23	(2010)
Total fertility rate (per woman)	7.0	(2011)
Adolescent birth rate (per 1000 women)	199	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 12 (2011)
Low birthweight incidence (moderate and severe, %) 27 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Nigeria


May 2013

DEMOGRAPHICS

Total population (000)	162,471	(2011)
Total under-five population (000)	27,215	(2011)
Births (000)	6,458	(2011)
Birth registration (%)	30	(2008)
Total under-five deaths (000)	756	(2011)
Neonatal deaths: % of all under-5 deaths	34	(2011)
Neonatal mortality rate (per 1000 live births)	39	(2011)
Infant mortality rate (per 1000 live births)	78	(2011)
Stillbirth rate (per 1000 total births)	42	(2009)
Total maternal deaths	40,000	(2010)
Lifetime risk of maternal death (1 in N)	29	(2010)
Total fertility rate (per woman)	5.5	(2011)
Adolescent birth rate (per 1000 women)	123	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %) 14 (2008)

Low birthweight incidence (moderate and severe, %) 12 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 38 (2008)

Introduction of solid, semi-solid/soft foods (%) 76 (2008)

Vitamin A two dose coverage (%) 73 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Pakistan


May 2013

DEMOGRAPHICS

Total population (000)	176,745	(2011)
Total under-five population (000)	22,113	(2011)
Births (000)	4,764	(2011)
Birth registration (%)	27	(2006-2007)
Total under-five deaths (000)	352	(2011)
Neonatal deaths: % of all under-5 deaths	48	(2011)
Neonatal mortality rate (per 1000 live births)	36	(2011)
Infant mortality rate (per 1000 live births)	59	(2011)
Stillbirth rate (per 1000 total births)	47	(2009)
Total maternal deaths	12,000	(2010)
Lifetime risk of maternal death (1 in N)	110	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	16	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2006-2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2011)
Low birthweight incidence (moderate and severe, %) 32 (2006-2007)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted




Note: Based on 2006 WHO reference population

Early initiation of breastfeeding (within 1 hr of birth, %) 29 (2006-2007)
Introduction of solid, semi-solid/soft foods (%) 36 (2006-2007)
Vitamin A two dose coverage (%) 90 (2011)

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Papua New Guinea


May 2013

DEMOGRAPHICS

Total population (000)	7,014	(2011)
Total under-five population (000)	975	(2011)
Births (000)	208	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	39	(2011)
Neonatal mortality rate (per 1000 live births)	23	(2011)
Infant mortality rate (per 1000 live births)	45	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	480	(2010)
Lifetime risk of maternal death (1 in N)	110	(2010)
Total fertility rate (per woman)	3.9	(2011)
Adolescent birth rate (per 1000 women)	70	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2005)
Low birthweight incidence (moderate and severe, %) 11 (2005)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Peru


May 2013

DEMOGRAPHICS

Total population (000)	29,400	(2011)
Total under-five population (000)	2,904	(2011)
Births (000)	591	(2011)
Birth registration (%)	93	(2007)
Total under-five deaths (000)	11	(2011)
Neonatal deaths: % of all under-5 deaths	51	(2011)
Neonatal mortality rate (per 1000 live births)	9	(2011)
Infant mortality rate (per 1000 live births)	14	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	400	(2010)
Lifetime risk of maternal death (1 in N)	570	(2010)
Total fertility rate (per woman)	2.5	(2011)
Adolescent birth rate (per 1000 women)	72	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2004


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 0 (2011)

Low birthweight incidence (moderate and severe, %) 8 (2007)


Early initiation of breastfeeding (within 1 hr of birth, %) 51 (2010)

Introduction of solid, semi-solid/soft foods (%) 82 (2011)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Philippines


May 2013

DEMOGRAPHICS

Total population (000)	94,852	(2011)
Total under-five population (000)	11,151	(2011)
Births (000)	2,358	(2011)
Birth registration (%)	83	(2008)
Total under-five deaths (000)	57	(2011)
Neonatal deaths: % of all under-5 deaths	50	(2011)
Neonatal mortality rate (per 1000 live births)	12	(2011)
Infant mortality rate (per 1000 live births)	20	(2011)
Stillbirth rate (per 1000 total births)	16	(2009)
Total maternal deaths	2,300	(2010)
Lifetime risk of maternal death (1 in N)	300	(2010)
Total fertility rate (per woman)	3.1	(2011)
Adolescent birth rate (per 1000 women)	53	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2008)

Low birthweight incidence (moderate and severe, %) 21 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 54 (2008)

Introduction of solid, semi-solid/soft foods (%) 90 (2008)

Vitamin A two dose coverage (%) 91 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Rwanda


May 2013

DEMOGRAPHICS

Total population (000)	10,943	(2011)
Total under-five population (000)	1,912	(2011)
Births (000)	449	(2011)
Birth registration (%)	63	(2010)
Total under-five deaths (000)	23	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	21	(2011)
Infant mortality rate (per 1000 live births)	38	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,500	(2010)
Lifetime risk of maternal death (1 in N)	54	(2010)
Total fertility rate (per woman)	5.3	(2011)
Adolescent birth rate (per 1000 women)	41	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2010


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 3 (2010)
Low birthweight incidence (moderate and severe, %) 7 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

São Tomé and Príncipe


May 2013

DEMOGRAPHICS

Total population (000)	169	(2011)
Total under-five population (000)	24	(2011)
Births (000)	5	(2011)
Birth registration (%)	75	(2008-2009)
Total under-five deaths (000)	0	(2011)
Neonatal deaths: % of all under-5 deaths	34	(2011)
Neonatal mortality rate (per 1000 live births)	29	(2011)
Infant mortality rate (per 1000 live births)	58	(2011)
Stillbirth rate (per 1000 total births)	22	(2009)
Total maternal deaths	4	(2010)
Lifetime risk of maternal death (1 in N)	330	(2010)
Total fertility rate (per woman)	3.6	(2011)
Adolescent birth rate (per 1000 women)	110	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2008-2009


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 11 (2008-2009)
Low birthweight incidence (moderate and severe, %) 8 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Senegal


May 2013

DEMOGRAPHICS

Total population (000)	12,768	(2011)
Total under-five population (000)	2,127	(2011)
Births (000)	471	(2011)
Birth registration (%)	75	(2010-2011)
Total under-five deaths (000)	30	(2011)
Neonatal deaths: % of all under-5 deaths	41	(2011)
Neonatal mortality rate (per 1000 live births)	26	(2011)
Infant mortality rate (per 1000 live births)	47	(2011)
Stillbirth rate (per 1000 total births)	34	(2009)
Total maternal deaths	1,700	(2010)
Lifetime risk of maternal death (1 in N)	54	(2010)
Total fertility rate (per woman)	4.7	(2011)
Adolescent birth rate (per 1000 women)	93	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2010-2011


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 10 (2010-2011)
Low birthweight incidence (moderate and severe, %) 19 (2010-2011)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted




Note: Based on 2006 WHO reference population

Early initiation of breastfeeding (within 1 hr of birth, %) 23 (2005)
Introduction of solid, semi-solid/soft foods (%) 61 (2010-2011)
Vitamin A two dose coverage (%) -

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Sierra Leone


May 2013

DEMOGRAPHICS

Total population (000)	5,997	(2011)
Total under-five population (000)	985	(2011)
Births (000)	227	(2011)
Birth registration (%)	78	(2010)
Total under-five deaths (000)	42	(2011)
Neonatal deaths: % of all under-5 deaths	27	(2011)
Neonatal mortality rate (per 1000 live births)	49	(2011)
Infant mortality rate (per 1000 live births)	119	(2011)
Stillbirth rate (per 1000 total births)	30	(2009)
Total maternal deaths	2,000	(2010)
Lifetime risk of maternal death (1 in N)	23	(2010)
Total fertility rate (per woman)	4.9	(2011)
Adolescent birth rate (per 1000 women)	98	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 9 (2010)

Low birthweight incidence (moderate and severe, %) 11 (2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 45 (2010)

Introduction of solid, semi-solid/soft foods (%) 25 (2008)

Vitamin A two dose coverage (%) 99 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Solomon Islands


May 2013

DEMOGRAPHICS

Total population (000)	552	(2011)
Total under-five population (000)	81	(2011)
Births (000)	17	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	0	(2011)
Neonatal deaths: % of all under-5 deaths	50	(2011)
Neonatal mortality rate (per 1000 live births)	11	(2011)
Infant mortality rate (per 1000 live births)	18	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	16	(2010)
Lifetime risk of maternal death (1 in N)	240	(2010)
Total fertility rate (per woman)	4.2	(2011)
Adolescent birth rate (per 1000 women)	70	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




Source: DHS

NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2007)
Low birthweight incidence (moderate and severe, %) 13 (2007)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted




Note: Based on 2006 WHO reference population

Early initiation of breastfeeding (within 1 hr of birth, %) 75 (2007)
Introduction of solid, semi-solid/soft foods (%) 81 (2007)
Vitamin A two dose coverage (%) -

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Somalia


May 2013

DEMOGRAPHICS

Total population (000)	9,557	(2011)
Total under-five population (000)	1,705	(2011)
Births (000)	416	(2011)
Birth registration (%)	3	(2006)
Total under-five deaths (000)	71	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	50	(2011)
Infant mortality rate (per 1000 live births)	108	(2011)
Stillbirth rate (per 1000 total births)	30	(2009)
Total maternal deaths	4,200	(2010)
Lifetime risk of maternal death (1 in N)	16	(2010)
Total fertility rate (per woman)	6.3	(2011)
Adolescent birth rate (per 1000 women)	123	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



MICS


NUTRITION

Wasting prevalence (moderate and severe, %) 13 (2006)
Low birthweight incidence (moderate and severe, %) -

Early initiation of breastfeeding (within 1 hr of birth, %) 26 (2006)
Introduction of solid, semi-solid/soft foods (%) 16 (2006)
Vitamin A two dose coverage (%) 12 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

South Africa


May 2013

DEMOGRAPHICS

Total population (000)	50,460	(2011)
Total under-five population (000)	4,982	(2011)
Births (000)	1,052	(2011)
Birth registration (%)	92	(2008)
Total under-five deaths (000)	47	(2011)
Neonatal deaths: % of all under-5 deaths	43	(2011)
Neonatal mortality rate (per 1000 live births)	19	(2011)
Infant mortality rate (per 1000 live births)	35	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	3,200	(2010)
Lifetime risk of maternal death (1 in N)	140	(2010)
Total fertility rate (per woman)	2.4	(2011)
Adolescent birth rate (per 1000 women)	54	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2003)

Low birthweight incidence (moderate and severe, %) -


Early initiation of breastfeeding (within 1 hr of birth, %) 61 (2003)

Introduction of solid, semi-solid/soft foods (%) 49 (2003)

Vitamin A two dose coverage (%) 44 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

South Sudan


May 2013

DEMOGRAPHICS

Total population (000)	10,314	(2011)
Total under-five population (000)	-	-
Births (000)	-	-
Birth registration (%)	35	(2010)
Total under-five deaths (000)	43	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	38	(2011)
Infant mortality rate (per 1000 live births)	76	(2011)
Stillbirth rate (per 1000 total births)	-	-
Total maternal deaths	-	-
Lifetime risk of maternal death (1 in N)	-	-
Total fertility rate (per woman)	-	-
Adolescent birth rate (per 1000 women)	-	-

Under-five mortality rate

Deaths per 1000 live births



Source: IGME 2012


Maternal mortality ratio

Deaths per 100,000 live births

No Data

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




MICS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



MICS

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



MICS

NUTRITION

Wasting prevalence (moderate and severe, %) 23 (2010)

Low birthweight incidence (moderate and severe, %) -


Early initiation of breastfeeding (within 1 hr of birth, %) -

Introduction of solid, semi-solid/soft foods (%) 21 (2010)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



MICS



Countdown to 2015

Maternal, Newborn & Child Survival

Sudan


May 2013

DEMOGRAPHICS

Total population (000)	34,318	(2011)
Total under-five population (000)	-	-
Births (000)	-	-
Birth registration (%)	59	(2010)
Total under-five deaths (000)	95	(2011)
Neonatal deaths: % of all under-5 deaths	37	(2011)
Neonatal mortality rate (per 1000 live births)	31	(2011)
Infant mortality rate (per 1000 live births)	57	(2011)
Stillbirth rate (per 1000 total births)	-	-
Total maternal deaths	-	-
Lifetime risk of maternal death (1 in N)	-	-
Total fertility rate (per woman)	-	-
Adolescent birth rate (per 1000 women)	-	-

Under-five mortality rate

Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio


Deaths per 100,000 live births

No Data

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH


Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 16 (2010)
Low birthweight incidence (moderate and severe, %) -

Early initiation of breastfeeding (within 1 hr of birth, %) -
Introduction of solid, semi-solid/soft foods (%) 51 (2010)
Vitamin A two dose coverage (%) -


Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Note: Based on 2006 WHO reference population



Countdown to 2015

Maternal, Newborn & Child Survival

Swaziland


May 2013

DEMOGRAPHICS

Total population (000)	1,203	(2011)
Total under-five population (000)	158	(2011)
Births (000)	35	(2011)
Birth registration (%)	50	(2010)
Total under-five deaths (000)	4	(2011)
Neonatal deaths: % of all under-5 deaths	34	(2011)
Neonatal mortality rate (per 1000 live births)	35	(2011)
Infant mortality rate (per 1000 live births)	69	(2011)
Stillbirth rate (per 1000 total births)	18	(2009)
Total maternal deaths	110	(2010)
Lifetime risk of maternal death (1 in N)	95	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	111	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 1 (2010)
Low birthweight incidence (moderate and severe, %) 9 (2010)

Early initiation of breastfeeding (within 1 hr of birth, %) 55 (2010)
Introduction of solid, semi-solid/soft foods (%) 66 (2010)
Vitamin A two dose coverage (%) 41 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Tajikistan


May 2013

DEMOGRAPHICS

Total population (000)	6,977	(2011)
Total under-five population (000)	883	(2011)
Births (000)	194	(2011)
Birth registration (%)	88	(2005)
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	40	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	12	(2009)
Total maternal deaths	130	(2010)
Lifetime risk of maternal death (1 in N)	430	(2010)
Total fertility rate (per woman)	3.2	(2011)
Adolescent birth rate (per 1000 women)	27	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2005

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2007)

Low birthweight incidence (moderate and severe, %) 10 (2005)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Tanzania, United Republic of


May 2013

DEMOGRAPHICS

Total population (000)	46,218	(2011)
Total under-five population (000)	8,273	(2011)
Births (000)	1,913	(2011)
Birth registration (%)	16	(2010)
Total under-five deaths (000)	122	(2011)
Neonatal deaths: % of all under-5 deaths	40	(2011)
Neonatal mortality rate (per 1000 live births)	25	(2011)
Infant mortality rate (per 1000 live births)	45	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	8,500	(2010)
Lifetime risk of maternal death (1 in N)	38	(2010)
Total fertility rate (per woman)	5.5	(2011)
Adolescent birth rate (per 1000 women)	128	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2010)
Low birthweight incidence (moderate and severe, %) 8 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Togo


May 2013

DEMOGRAPHICS

Total population (000)	6,155	(2011)
Total under-five population (000)	870	(2011)
Births (000)	195	(2011)
Birth registration (%)	78	(2010)
Total under-five deaths (000)	21	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	36	(2011)
Infant mortality rate (per 1000 live births)	73	(2011)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	580	(2010)
Lifetime risk of maternal death (1 in N)	80	(2010)
Total fertility rate (per woman)	4.0	(2011)
Adolescent birth rate (per 1000 women)	89	(1995)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2010)
Low birthweight incidence (moderate and severe, %) 11 (2010)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Turkmenistan


May 2013

DEMOGRAPHICS

Total population (000)	5,105	(2011)
Total under-five population (000)	499	(2011)
Births (000)	109	(2011)
Birth registration (%)	96	(2006)
Total under-five deaths (000)	5	(2011)
Neonatal deaths: % of all under-5 deaths	44	(2011)
Neonatal mortality rate (per 1000 live births)	22	(2011)
Infant mortality rate (per 1000 live births)	45	(2011)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	73	(2010)
Lifetime risk of maternal death (1 in N)	590	(2010)
Total fertility rate (per woman)	2.4	(2011)
Adolescent birth rate (per 1000 women)	21	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate

No Data

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



No Data


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2006)
Low birthweight incidence (moderate and severe, %) 4 (2006)

Early initiation of breastfeeding (within 1 hr of birth, %) 60 (2006)
Introduction of solid, semi-solid/soft foods (%) 54 (2005)
Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Uganda


May 2013

DEMOGRAPHICS

Total population (000)	34,509	(2011)
Total under-five population (000)	6,643	(2011)
Births (000)	1,545	(2011)
Birth registration (%)	30	(2011)
Total under-five deaths (000)	131	(2011)
Neonatal deaths: % of all under-5 deaths	33	(2011)
Neonatal mortality rate (per 1000 live births)	28	(2011)
Infant mortality rate (per 1000 live births)	58	(2011)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	4,700	(2010)
Lifetime risk of maternal death (1 in N)	49	(2010)
Total fertility rate (per woman)	6.1	(2011)
Adolescent birth rate (per 1000 women)	159	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2011

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2011)
Low birthweight incidence (moderate and severe, %) 14 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Uzbekistan


May 2013

DEMOGRAPHICS

Total population (000)	27,760	(2011)
Total under-five population (000)	2,806	(2011)
Births (000)	589	(2011)
Birth registration (%)	100	(2009)
Total under-five deaths (000)	30	(2011)
Neonatal deaths: % of all under-5 deaths	30	(2011)
Neonatal mortality rate (per 1000 live births)	15	(2011)
Infant mortality rate (per 1000 live births)	42	(2011)
Stillbirth rate (per 1000 total births)	6	(2009)
Total maternal deaths	160	(2010)
Lifetime risk of maternal death (1 in N)	1,400	(2010)
Total fertility rate (per woman)	2.3	(2011)
Adolescent birth rate (per 1000 women)	26	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2006)

Low birthweight incidence (moderate and severe, %) 5 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 67 (2006)

Introduction of solid, semi-solid/soft foods (%) 47 (2006)

Vitamin A two dose coverage (%) 95 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Vietnam


May 2013

DEMOGRAPHICS

Total population (000)	88,792	(2011)
Total under-five population (000)	7,202	(2011)
Births (000)	1,458	(2011)
Birth registration (%)	95	(2011)
Total under-five deaths (000)	32	(2011)
Neonatal deaths: % of all under-5 deaths	55	(2011)
Neonatal mortality rate (per 1000 live births)	12	(2011)
Infant mortality rate (per 1000 live births)	17	(2011)
Stillbirth rate (per 1000 total births)	13	(2009)
Total maternal deaths	860	(2010)
Lifetime risk of maternal death (1 in N)	870	(2010)
Total fertility rate (per woman)	1.8	(2011)
Adolescent birth rate (per 1000 women)	35	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2011


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2011)
Low birthweight incidence (moderate and severe, %) 5 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Yemen


May 2013

DEMOGRAPHICS

Total population (000)	24,800	(2011)
Total under-five population (000)	4,181	(2011)
Births (000)	940	(2011)
Birth registration (%)	22	(2006)
Total under-five deaths (000)	70	(2011)
Neonatal deaths: % of all under-5 deaths	43	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	57	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,900	(2010)
Lifetime risk of maternal death (1 in N)	90	(2010)
Total fertility rate (per woman)	5.1	(2011)
Adolescent birth rate (per 1000 women)	80	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)


■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate



Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2003)
Low birthweight incidence (moderate and severe, %) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Zambia


May 2013

DEMOGRAPHICS

Total population (000)	13,475	(2011)
Total under-five population (000)	2,510	(2011)
Births (000)	622	(2011)
Birth registration (%)	14	(2007)
Total under-five deaths (000)	46	(2011)
Neonatal deaths: % of all under-5 deaths	37	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	26	(2009)
Total maternal deaths	2,600	(2010)
Lifetime risk of maternal death (1 in N)	37	(2010)
Total fertility rate (per woman)	6.3	(2011)
Adolescent birth rate (per 1000 women)	151	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2007)
Low birthweight incidence (moderate and severe, %) 11 (2007)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Zimbabwe


May 2013

DEMOGRAPHICS

Total population (000)	12,754	(2011)
Total under-five population (000)	1,719	(2011)
Births (000)	377	(2011)
Birth registration (%)	49	(2010-2011)
Total under-five deaths (000)	24	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	30	(2011)
Infant mortality rate (per 1000 live births)	43	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	2,200	(2010)
Lifetime risk of maternal death (1 in N)	52	(2010)
Total fertility rate (per woman)	3.2	(2011)
Adolescent birth rate (per 1000 women)	115	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2010-2011


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 3 (2010-2011)

Low birthweight incidence (moderate and severe, %) 11 (2005-2006)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Early initiation of breastfeeding (within 1 hr of birth, %) 69 (2005-2006)

Introduction of solid, semi-solid/soft foods (%) 86 (2010-2011)

Vitamin A two dose coverage (%) 56 (2011)

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

