



Country profiles

The information in the country profiles, customized to showcase the core indicators selected by the Commission on Information and Accountability for Women's and Children's Health, is intended to help policy makers and their partners assess progress and prioritize actions to save women's and children's lives. These country profiles reflect the data that are available for each country. Most of the data for the Commission indicators come from household surveys: mainly the USAID-supported Demographic and Health Surveys and the UNICEF-supported Multiple Indicator Cluster Surveys.^{xxix} More information on the data sources for the Commission indicators is available in the annexes to previous Countdown reports: www.countdown2015mnch.org/reports-and-articles.

New survey data are available for 22 countries since the publication of Countdown's 2012 Report. Coverage estimates that are missing or that are more than five years old indicate an urgent need for concerted action to increase data collection efforts so that updated evidence is available for policy and program development.

This report contains profiles for 75 countries with high burdens of maternal and/or child mortality, including the 49 lowest-income countries covered by the Global Strategy for Women's and Children's Health. Profiles are included for the following countries:

Afghanistan	Gambia	Nigeria
Angola	Ghana	Pakistan
Azerbaijan	Guatemala	Papua New Guinea
Bangladesh	Guinea	Peru
Benin	Guinea-Bissau	Philippines
Bolivia	Haiti	Rwanda
Botswana	India	São Tomé and Príncipe
Brazil	Indonesia	Senegal
Burkina Faso	Iraq	Sierra Leone
Burundi	Kenya	Solomon Islands
Cambodia	Korea, Democratic People's Republic of	Somalia
Cameroon	Kyrgyzstan	South Africa
Central African Republic	Lao People's Democratic Republic	South Sudan
Chad	Lesotho	Sudan
China	Liberia	Swaziland
Comoros	Madagascar	Tajikistan
Congo	Malawi	Tanzania, United Republic of
Congo, Democratic Republic of the	Mali	Togo
Côte d'Ivoire	Mauritania	Turkmenistan
Djibouti	Mexico	Uganda
Egypt	Morocco	Uzbekistan
Equatorial Guinea	Mozambique	Vietnam
Eritrea	Myanmar	Yemen
Ethiopia	Nepal	Zambia
Gabon	Niger	Zimbabwe



Countdown to 2015

Maternal, Newborn & Child Survival

Lao People's Democratic Republic


May 2013

DEMOGRAPHICS

Total population (000)	6,288	(2011)
Total under-five population (000)	682	(2011)
Births (000)	140	(2011)
Birth registration (%)	72	(2006)
Total under-five deaths (000)	6	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	18	(2011)
Infant mortality rate (per 1000 live births)	34	(2011)
Stillbirth rate (per 1000 total births)	14	(2009)
Total maternal deaths	670	(2010)
Lifetime risk of maternal death (1 in N)	74	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	110	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2006)
Low birthweight incidence (moderate and severe, %) 11 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Lesotho


May 2013

DEMOGRAPHICS

Total population (000)	2,194	(2011)
Total under-five population (000)	276	(2011)
Births (000)	60	(2011)
Birth registration (%)	45	(2009)
Total under-five deaths (000)	5	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	39	(2011)
Infant mortality rate (per 1000 live births)	63	(2011)
Stillbirth rate (per 1000 total births)	25	(2009)
Total maternal deaths	370	(2010)
Lifetime risk of maternal death (1 in N)	53	(2010)
Total fertility rate (per woman)	3.1	(2011)
Adolescent birth rate (per 1000 women)	92	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS, DHS

NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2009)

Low birthweight incidence (moderate and severe, %) 11 (2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 53 (2009)

Introduction of solid, semi-solid/soft foods (%) 68 (2009)

Vitamin A two dose coverage (%) -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: MICS, DHS

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Liberia


May 2013

DEMOGRAPHICS

Total population (000)	4,129	(2011)
Total under-five population (000)	696	(2011)
Births (000)	157	(2011)
Birth registration (%)	4	(2007)
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	36	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	58	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	24	(2010)
Total fertility rate (per woman)	5.2	(2011)
Adolescent birth rate (per 1000 women)	177	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %)	3	(2010)
Low birthweight incidence (moderate and severe, %)	14	(2007)

Early initiation of breastfeeding (within 1 hr of birth, %)	44	(2010)
Introduction of solid, semi-solid/soft foods (%)	51	(2010)
Vitamin A two dose coverage (%)	96	(2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Madagascar


May 2013

DEMOGRAPHICS

Total population (000)	21,315	(2011)
Total under-five population (000)	3,379	(2011)
Births (000)	747	(2011)
Birth registration (%)	80	(2008-2009)
Total under-five deaths (000)	45	(2011)
Neonatal deaths: % of all under-5 deaths	38	(2011)
Neonatal mortality rate (per 1000 live births)	23	(2011)
Infant mortality rate (per 1000 live births)	43	(2011)
Stillbirth rate (per 1000 total births)	21	(2009)
Total maternal deaths	1,800	(2010)
Lifetime risk of maternal death (1 in N)	81	(2010)
Total fertility rate (per woman)	4.6	(2011)
Adolescent birth rate (per 1000 women)	147	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008-2009

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2003-2004)

Low birthweight incidence (moderate and severe, %) 16 (2008-2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 72 (2008-2009)

Introduction of solid, semi-solid/soft foods (%) 86 (2008-2009)

Vitamin A two dose coverage (%) 91 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Malawi


May 2013

DEMOGRAPHICS

Total population (000)	15,381	(2011)
Total under-five population (000)	2,832	(2011)
Births (000)	686	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	52	(2011)
Neonatal deaths: % of all under-5 deaths	35	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	53	(2011)
Stillbirth rate (per 1000 total births)	24	(2009)
Total maternal deaths	3,000	(2010)
Lifetime risk of maternal death (1 in N)	36	(2010)
Total fertility rate (per woman)	6.0	(2011)
Adolescent birth rate (per 1000 women)	157	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2010

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 4 (2010)

Low birthweight incidence (moderate and severe, %) 13 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 58 (2006)

Introduction of solid, semi-solid/soft foods (%) 86 (2010)

Vitamin A two dose coverage (%) 96 (2011)


Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Note: Based on 2006 WHO reference population

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mali


May 2013

DEMOGRAPHICS

Total population (000)	15,840	(2011)
Total under-five population (000)	2,998	(2011)
Births (000)	728	(2011)
Birth registration (%)	81	(2010)
Total under-five deaths (000)	121	(2011)
Neonatal deaths: % of all under-5 deaths	29	(2011)
Neonatal mortality rate (per 1000 live births)	49	(2011)
Infant mortality rate (per 1000 live births)	98	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	3,800	(2010)
Lifetime risk of maternal death (1 in N)	28	(2010)
Total fertility rate (per woman)	6.2	(2011)
Adolescent birth rate (per 1000 women)	190	(2004)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

Percent HIV+ pregnant women receiving ARVs for PMTCT

Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: Poorest 20% Richest 20%




DHS 2006

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


Percent of children immunized against measles
Percent of children immunized with 3 doses DTP
Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

Percent children <5 years with suspected pneumonia taken to appropriate health provider
Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

15

(2006)

Low birthweight incidence (moderate and severe, %)

19

(2006)

Early initiation of breastfeeding (within 1 hr of birth, %)

46

(2006)

Introduction of solid, semi-solid/soft foods (%)

25

(2006)


Vitamin A two dose coverage (%)

96

(2011)

Underweight and stunting prevalence


Percent children <5 years who are underweight
Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Mauritania


May 2013

DEMOGRAPHICS

Total population (000)	3,542	(2011)
Total under-five population (000)	522	(2011)
Births (000)	118	(2011)
Birth registration (%)	56	(2007)
Total under-five deaths (000)	13	(2011)
Neonatal deaths: % of all under-5 deaths	37	(2011)
Neonatal mortality rate (per 1000 live births)	40	(2011)
Infant mortality rate (per 1000 live births)	76	(2011)
Stillbirth rate (per 1000 total births)	27	(2009)
Total maternal deaths	590	(2010)
Lifetime risk of maternal death (1 in N)	44	(2010)
Total fertility rate (per woman)	4.5	(2011)
Adolescent birth rate (per 1000 women)	88	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



MICS 2007


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 12 (2010)
Low birthweight incidence (moderate and severe, %) 34 (2007)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Mexico


May 2013

DEMOGRAPHICS

Total population (000)	114,793	(2011)
Total under-five population (000)	10,938	(2011)
Births (000)	2,195	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	34	(2011)
Neonatal deaths: % of all under-5 deaths	46	(2011)
Neonatal mortality rate (per 1000 live births)	7	(2011)
Infant mortality rate (per 1000 live births)	13	(2011)
Stillbirth rate (per 1000 total births)	5	(2009)
Total maternal deaths	1,100	(2010)
Lifetime risk of maternal death (1 in N)	790	(2010)
Total fertility rate (per woman)	2.3	(2011)
Adolescent birth rate (per 1000 women)	87	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
□ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: UNICEF/UNAIDS/WHO

NUTRITION

Wasting prevalence (moderate and severe, %) - -

Low birthweight incidence (moderate and severe, %) 7 (2009)


Early initiation of breastfeeding (within 1 hr of birth, %) 18 (2009)

Introduction of solid, semi-solid/soft foods (%) 27 (2009)

Vitamin A two dose coverage (%) - -

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Morocco


May 2013

DEMOGRAPHICS

Total population (000)	32,273	(2011)
Total under-five population (000)	3,051	(2011)
Births (000)	620	(2011)
Birth registration (%)	85	(2000)
Total under-five deaths (000)	21	(2011)
Neonatal deaths: % of all under-5 deaths	56	(2011)
Neonatal mortality rate (per 1000 live births)	19	(2011)
Infant mortality rate (per 1000 live births)	28	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	650	(2010)
Lifetime risk of maternal death (1 in N)	400	(2010)
Total fertility rate (per woman)	2.2	(2011)
Adolescent birth rate (per 1000 women)	18	(2005)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2003-2004


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %)

2 (2010-2011)

Low birthweight incidence (moderate and severe, %)

15 (2003-2004)

Early initiation of breastfeeding (within 1 hr of birth, %)

52 (2003-2004)

Introduction of solid, semi-solid/soft foods (%)


66 (2003-2004)

Vitamin A two dose coverage (%)

-

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Mozambique


May 2013

DEMOGRAPHICS

Total population (000)	23,930	(2011)
Total under-five population (000)	3,875	(2011)
Births (000)	889	(2011)
Birth registration (%)	31	(2008)
Total under-five deaths (000)	86	(2011)
Neonatal deaths: % of all under-5 deaths	35	(2011)
Neonatal mortality rate (per 1000 live births)	34	(2011)
Infant mortality rate (per 1000 live births)	72	(2011)
Stillbirth rate (per 1000 total births)	28	(2009)
Total maternal deaths	4,300	(2010)
Lifetime risk of maternal death (1 in N)	43	(2010)
Total fertility rate (per woman)	4.8	(2011)
Adolescent birth rate (per 1000 women)	193	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




MICS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 6 (2011)
Low birthweight incidence (moderate and severe, %) 16 (2008)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Myanmar


May 2013

DEMOGRAPHICS

Total population (000)	48,337	(2011)
Total under-five population (000)	3,985	(2011)
Births (000)	824	(2011)
Birth registration (%)	72	(2009-2010)
Total under-five deaths (000)	53	(2011)
Neonatal deaths: % of all under-5 deaths	47	(2011)
Neonatal mortality rate (per 1000 live births)	30	(2011)
Infant mortality rate (per 1000 live births)	48	(2011)
Stillbirth rate (per 1000 total births)	20	(2009)
Total maternal deaths	1,600	(2010)
Lifetime risk of maternal death (1 in N)	250	(2010)
Total fertility rate (per woman)	2.0	(2011)
Adolescent birth rate (per 1000 women)	17	(2001)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery

Percent live births attended by skilled health personnel




Source: DHS, MICS, Other NS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



Source: MICS

NUTRITION

Wasting prevalence (moderate and severe, %) 8 (2009-2010)

Low birthweight incidence (moderate and severe, %) 9 (2009-2010)


Early initiation of breastfeeding (within 1 hr of birth, %) 76 (2009-2010)

Introduction of solid, semi-solid/soft foods (%) 81 (2009-2010)

Vitamin A two dose coverage (%) 96 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Source: MICS



Countdown to 2015

Maternal, Newborn & Child Survival

Nepal


May 2013

DEMOGRAPHICS

Total population (000)	30,486	(2011)
Total under-five population (000)	3,450	(2011)
Births (000)	722	(2011)
Birth registration (%)	42	(2011)
Total under-five deaths (000)	34	(2011)
Neonatal deaths: % of all under-5 deaths	58	(2011)
Neonatal mortality rate (per 1000 live births)	27	(2011)
Infant mortality rate (per 1000 live births)	39	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,200	(2010)
Lifetime risk of maternal death (1 in N)	190	(2010)
Total fertility rate (per woman)	2.7	(2011)
Adolescent birth rate (per 1000 women)	81	(2010)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2011

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 11 (2011)
Low birthweight incidence (moderate and severe, %) 18 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Niger


May 2013

DEMOGRAPHICS

Total population (000)	16,069	(2011)
Total under-five population (000)	3,200	(2011)
Births (000)	777	(2011)
Birth registration (%)	32	(2006)
Total under-five deaths (000)	89	(2011)
Neonatal deaths: % of all under-5 deaths	28	(2011)
Neonatal mortality rate (per 1000 live births)	32	(2011)
Infant mortality rate (per 1000 live births)	66	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	4,500	(2010)
Lifetime risk of maternal death (1 in N)	23	(2010)
Total fertility rate (per woman)	7.0	(2011)
Adolescent birth rate (per 1000 women)	199	(2003)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2006


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 12 (2011)
Low birthweight incidence (moderate and severe, %) 27 (2006)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Nigeria


May 2013

DEMOGRAPHICS

Total population (000)	162,471	(2011)
Total under-five population (000)	27,215	(2011)
Births (000)	6,458	(2011)
Birth registration (%)	30	(2008)
Total under-five deaths (000)	756	(2011)
Neonatal deaths: % of all under-5 deaths	34	(2011)
Neonatal mortality rate (per 1000 live births)	39	(2011)
Infant mortality rate (per 1000 live births)	78	(2011)
Stillbirth rate (per 1000 total births)	42	(2009)
Total maternal deaths	40,000	(2010)
Lifetime risk of maternal death (1 in N)	29	(2010)
Total fertility rate (per woman)	5.5	(2011)
Adolescent birth rate (per 1000 women)	123	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Source: DHS

Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH


Immunization



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 14 (2008)
Low birthweight incidence (moderate and severe, %) 12 (2008)

Early initiation of breastfeeding (within 1 hr of birth, %) 38 (2008)
Introduction of solid, semi-solid/soft foods (%) 76 (2008)
Vitamin A two dose coverage (%) 73 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Pakistan


May 2013

DEMOGRAPHICS

Total population (000)	176,745	(2011)
Total under-five population (000)	22,113	(2011)
Births (000)	4,764	(2011)
Birth registration (%)	27	(2006-2007)
Total under-five deaths (000)	352	(2011)
Neonatal deaths: % of all under-5 deaths	48	(2011)
Neonatal mortality rate (per 1000 live births)	36	(2011)
Infant mortality rate (per 1000 live births)	59	(2011)
Stillbirth rate (per 1000 total births)	47	(2009)
Total maternal deaths	12,000	(2010)
Lifetime risk of maternal death (1 in N)	110	(2010)
Total fertility rate (per woman)	3.3	(2011)
Adolescent birth rate (per 1000 women)	16	(2007)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2006-2007

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 15 (2011)
Low birthweight incidence (moderate and severe, %) 32 (2006-2007)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted




Note: Based on 2006 WHO reference population

Early initiation of breastfeeding (within 1 hr of birth, %) 29 (2006-2007)
Introduction of solid, semi-solid/soft foods (%) 36 (2006-2007)
Vitamin A two dose coverage (%) 90 (2011)

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Papua New Guinea


May 2013

DEMOGRAPHICS

Total population (000)	7,014	(2011)
Total under-five population (000)	975	(2011)
Births (000)	208	(2011)
Birth registration (%)	-	-
Total under-five deaths (000)	12	(2011)
Neonatal deaths: % of all under-5 deaths	39	(2011)
Neonatal mortality rate (per 1000 live births)	23	(2011)
Infant mortality rate (per 1000 live births)	45	(2011)
Stillbirth rate (per 1000 total births)	15	(2009)
Total maternal deaths	480	(2010)
Lifetime risk of maternal death (1 in N)	110	(2010)
Total fertility rate (per woman)	3.9	(2011)
Adolescent birth rate (per 1000 women)	70	(2000)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 5 (2005)
Low birthweight incidence (moderate and severe, %) 11 (2005)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Peru


May 2013

DEMOGRAPHICS

Total population (000)	29,400	(2011)
Total under-five population (000)	2,904	(2011)
Births (000)	591	(2011)
Birth registration (%)	93	(2007)
Total under-five deaths (000)	11	(2011)
Neonatal deaths: % of all under-5 deaths	51	(2011)
Neonatal mortality rate (per 1000 live births)	9	(2011)
Infant mortality rate (per 1000 live births)	14	(2011)
Stillbirth rate (per 1000 total births)	10	(2009)
Total maternal deaths	400	(2010)
Lifetime risk of maternal death (1 in N)	570	(2010)
Total fertility rate (per woman)	2.5	(2011)
Adolescent birth rate (per 1000 women)	72	(2008)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2004


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 0 (2011)
Low birthweight incidence (moderate and severe, %) 8 (2007)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed





Countdown to 2015

Maternal, Newborn & Child Survival

Philippines


May 2013

DEMOGRAPHICS

Total population (000)	94,852	(2011)
Total under-five population (000)	11,151	(2011)
Births (000)	2,358	(2011)
Birth registration (%)	83	(2008)
Total under-five deaths (000)	57	(2011)
Neonatal deaths: % of all under-5 deaths	50	(2011)
Neonatal mortality rate (per 1000 live births)	12	(2011)
Infant mortality rate (per 1000 live births)	20	(2011)
Stillbirth rate (per 1000 total births)	16	(2009)
Total maternal deaths	2,300	(2010)
Lifetime risk of maternal death (1 in N)	300	(2010)
Total fertility rate (per woman)	3.1	(2011)
Adolescent birth rate (per 1000 women)	53	(2006)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




DHS 2008

Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

CHILD HEALTH

Immunization


■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics



NUTRITION

Wasting prevalence (moderate and severe, %) 7 (2008)

Low birthweight incidence (moderate and severe, %) 21 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 54 (2008)

Introduction of solid, semi-solid/soft foods (%) 90 (2008)

Vitamin A two dose coverage (%) 91 (2011)

Underweight and stunting prevalence


■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed



Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health



Countdown to 2015

Maternal, Newborn & Child Survival

Rwanda


May 2013

DEMOGRAPHICS

Total population (000)	10,943	(2011)
Total under-five population (000)	1,912	(2011)
Births (000)	449	(2011)
Birth registration (%)	63	(2010)
Total under-five deaths (000)	23	(2011)
Neonatal deaths: % of all under-5 deaths	42	(2011)
Neonatal mortality rate (per 1000 live births)	21	(2011)
Infant mortality rate (per 1000 live births)	38	(2011)
Stillbirth rate (per 1000 total births)	23	(2009)
Total maternal deaths	1,500	(2010)
Lifetime risk of maternal death (1 in N)	54	(2010)
Total fertility rate (per woman)	5.3	(2011)
Adolescent birth rate (per 1000 women)	41	(2009)

Under-five mortality rate


Deaths per 1000 live births



Source: IGME 2012

Maternal mortality ratio

Deaths per 100,000 live births




Source: MMEIG 2012

Note: MDG target calculated by Countdown to 2015.

MATERNAL AND NEWBORN HEALTH

Coverage along the continuum of care




Source: DHS, MICS, Other NS

* See Annex/website for indicator definition

Skilled attendant at delivery


Percent live births attended by skilled health personnel



Prevention of mother-to-child transmission of HIV

Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs)

■ Percent HIV+ pregnant women receiving ARVs for PMTCT
■ Uncertainty range around the estimate




Source: UNICEF/UNAIDS/WHO

EQUITY

Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%



DHS 2010


Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

Developed by Countdown to report on core indicators identified by the UN Commission on Information and Accountability, in support of the Global Strategy for Women's and Children's Health

CHILD HEALTH

Immunization

■ Percent of children immunized against measles
■ Percent of children immunized with 3 doses DTP
■ Percent of children immunized with 3 doses Hib



Source: WHO/UNICEF

Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider
■ Percent children <5 years with suspected pneumonia receiving antibiotics




NUTRITION

Wasting prevalence (moderate and severe, %) 3 (2010)
Low birthweight incidence (moderate and severe, %) 7 (2010)

Underweight and stunting prevalence

■ Percent children <5 years who are underweight
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population

Exclusive breastfeeding

Percent infants <6 months exclusively breastfed

